

Colombeau solutions to nonlinear wave equations

Michael Oberguggenberger*

Institut für Grundlagen der Baingenieurwissenschaften

Universität Innsbruck, A-6020 Innsbruck, Austria

E-mail: michael.oberguggenberger@uibk.ac.at

Abstract

This paper is a tutorial that demonstrates various methods from the Colombeau theory of generalized functions in the context of semilinear wave equations. The Colombeau generalized functions constitute differential algebras that contain the space of distributions. We solve the 1D-semilinear wave equation in these algebras, show how delta waves can be computed and then turn to linear and nonlinear regularity theory.

Keywords: Colombeau algebras, nonlinear wave equations, delta waves, regularity of solutions.

AMS Subject Classification: 35D05, 35L60, 46F30

1 Introduction

This tutorial serves the purpose of demonstrating how the theory of Colombeau algebras can be used to solve and study nonlinear partial differential equations that do not have solutions in the sense of distributions. It starts out with existence results obtained in the eighties of the twentieth century and leads up to recent regularity results.

For the purpose of exposition we focus on a model problem, the one-dimensional semilinear wave equation with singular initial data

$$\begin{aligned} \partial_t^2 u(x, t) - \partial_x^2 u(x, t) &= f(u(x, t)) + h(x, t), \quad (x, t) \in \mathbb{R}^2, \\ u(x, 0) &= a(x), \quad \partial_t u(x, 0) = b(x), \quad x \in \mathbb{R}, \end{aligned} \quad (1)$$

where f is a smooth, polynomially bounded function and a, b and h are Colombeau generalized functions on the real line and on the (x, t) -plane, respectively. The main questions to be addressed are the following:

- (a) existence and uniqueness of generalized solutions in Colombeau algebras;
- (b) limiting behavior of the representatives when the data are distributions;
- (c) regularity of generalized solutions.

Concerning question (a), a short survey of existence and uniqueness results can be found in [28]. The existence of Colombeau solutions to semilinear hyperbolic systems was one of the early applications of the theory [3, 23]. For linear hyperbolic systems with non-smooth (Colombeau generalized) coefficients,

*Partially supported by FWF(Austria), grant Y237 and by FAPESP (Brasil)

existence and uniqueness was established in [24], for symmetric hyperbolic systems in higher space dimensions in [18] and for hyperbolic pseudodifferential systems with Colombeau symbols in [15]. Notably for semilinear hyperbolic systems, question (b) has been answered in many cases, involving the notion of delta-waves [8, 13, 22, 26, 33, 34, 37]. Finally, regularity theory for Colombeau solutions is now based on the subalgebra $\mathcal{G}^\infty$ of regular Colombeau functions and currently an active area of research, making use of pseudodifferential and microlocal techniques [9, 11, 17, 20]. In particular, the propagation of the $\mathcal{G}^\infty$ -wave front set in linear systems with Colombeau coefficients is a theme of recent investigations [12]. To date, only few results are available about regularity of Colombeau solutions in the nonlinear case.

The plan of the tutorial is as follows. After recalling the required notions from Colombeau theory in Section 2, we shall prove existence and uniqueness of a solution u belonging to the Colombeau algebra on $\mathbb{R}^2$ in Section 3 and state various preliminary regularity properties. Section 4 is devoted to computing the associated distribution (the distributional limit of the representing nets) when the initial data are delta functions or derivatives of delta functions. In Section 5 we turn to regularity theory. We recall the $\mathcal{G}^\infty$ -regularity result for the linear case and show that it fails in the nonlinear case. The tutorial finishes with a recent result on propagation of regularity in the nonlinear case in Section 5.

For simplicity of presentation, we restrict our attention to the one-dimensional case and Lipschitz-continuous nonlinearity f . At the appropriate places of the tutorial, we will indicate what is known about the non-Lipschitz and the higher dimensional case. With Lipschitz-continuous f , in particular, the solution to (1) exists globally in space and time. Therefore, we need not enter the discussion of domains of existence nor use other versions of Colombeau algebras to accommodate energy estimates when f grows polynomially.

2 Notation

The paper is placed in the framework of algebras of generalized functions introduced by Colombeau in [6, 7]. We shall fix the notation and introduce a number of known as well as new classes of generalized functions here. For more details, see [14].

Let Ω be an open subset of $\mathbb{R}^n$. The basic objects of the theory as we use it are families $(u_\varepsilon)_{\varepsilon \in (0,1]}$ of smooth functions $u_\varepsilon \in \mathcal{C}^\infty(\Omega)$ for $0 < \varepsilon \leq 1$. We single out the following subalgebras:

Moderate families, denoted by $\mathcal{E}_M(\Omega)$, are defined by the property:

$$\forall K \Subset \Omega \forall \alpha \in \mathbb{N}_0^n \exists p \geq 0 : \sup_{x \in K} |\partial^\alpha u_\varepsilon(x)| = \mathcal{O}(\varepsilon^{-p}) \text{ as } \varepsilon \rightarrow 0. \quad (2)$$

Null families, denoted by $\mathcal{N}(\Omega)$, are defined by the property:

$$\forall K \Subset \Omega \forall \alpha \in \mathbb{N}_0^n \forall q \geq 0 : \sup_{x \in K} |\partial^\alpha u_\varepsilon(x)| = \mathcal{O}(\varepsilon^q) \text{ as } \varepsilon \rightarrow 0. \quad (3)$$

Thus moderate families satisfy a locally uniform polynomial estimate as $\varepsilon \rightarrow 0$, together with all derivatives, while null functionals vanish faster than any power

of ε in the same situation. The null families form a differential ideal in the collection of moderate families. The *Colombeau algebra* is the factor algebra

$$\mathcal{G}(\Omega) = \mathcal{E}_M(\Omega)/\mathcal{N}(\Omega).$$

The algebra $\mathcal{G}(\Omega)$ just defined coincides with the *special Colombeau algebra* in [14, Def. 1.2.2], where the notation $\mathcal{G}^s(\Omega)$ has been employed. It was called the *simplified Colombeau algebra* in [4].

Families $(r_\varepsilon)_{\varepsilon \in (0,1]}$ of complex numbers such that $|r_\varepsilon| = \mathcal{O}(\varepsilon^{-p})$ as $\varepsilon \rightarrow 0$ for some $p \geq 0$ are called *moderate*, those for which $|r_\varepsilon| = \mathcal{O}(\varepsilon^q)$ for every $q \geq 0$ are termed *negligible*. The ring $\tilde{\mathbb{C}}$ of Colombeau generalized numbers is obtained by factoring moderate families of complex numbers with respect to negligible families. When Ω is connected, $\tilde{\mathbb{C}}$ coincides with the *ring of constants* in the differential algebra $\mathcal{G}(\Omega)$.

The restriction of an element $u \in \mathcal{G}(\mathbb{R}^2)$ to the line $\{t = 0\}$ is defined on representatives by

$$u|_{\{t=0\}} = \text{class of } (u_\varepsilon(\cdot, 0))_{\varepsilon \in (0,1]}.$$

Similarly, restrictions of the elements of $\mathcal{G}(\Omega)$ to open subsets of Ω are defined on representatives. One can see that $\Omega \rightarrow \mathcal{G}(\Omega)$ is a sheaf of differential algebras on $\mathbb{R}^n$. The space of compactly supported distributions is imbedded in $\mathcal{G}(\Omega)$ by convolution:

$$\iota : \mathcal{E}'(\Omega) \rightarrow \mathcal{G}(\Omega), \quad \iota(w) = \text{class of } (w * (\varphi_\varepsilon)|_\Omega)_{\varepsilon \in (0,1]}, \quad (4)$$

where

$$\varphi_\varepsilon(x) = \varepsilon^{-n} \varphi(x/\varepsilon) \quad (5)$$

is obtained by scaling a fixed test function $\varphi \in \mathcal{S}(\mathbb{R}^n)$ of integral one with all moments vanishing. By the sheaf property, this can be extended in a unique way to an imbedding of the space of distributions $\mathcal{D}'(\Omega)$. In the case $\Omega = \mathbb{R}^n$, the imbedding is given by the explicit formula

$$\iota : \mathcal{D}'(\Omega) \rightarrow \mathcal{G}(\mathbb{R}^n), \quad \iota(w) = \text{class of } (w * (\chi \varphi_\varepsilon))_{\varepsilon \in (0,1]}, \quad (6)$$

where χ is some compactly supported smooth function identically equal to one in a neighborhood of zero. We refer to [14, Sec. 1.2] for further explicit expressions in general and in special cases.

One of the main features of the Colombeau construction is the fact that this imbedding renders $\mathcal{C}^\infty(\Omega)$ a faithful subalgebra. In fact, given $f \in \mathcal{C}^\infty(\Omega)$, one can define a corresponding element of $\mathcal{G}(\Omega)$ by the constant imbedding $\sigma(f) = \text{class of } [(\varepsilon, x) \rightarrow f(x)]$. Then the important equality $\iota(f) = \sigma(f)$ holds in $\mathcal{G}(\Omega)$.

If $u \in \mathcal{G}(\Omega)$ and f is a smooth function which is of at most polynomial growth at infinity, together with all its derivatives, the superposition $f(u)$ is a well-defined element of $\mathcal{G}(\Omega)$.

The algebra $\mathcal{G}(\Omega)$ can be equipped with a topology that turns it into a topological ring and a topological module over $\tilde{\mathbb{C}}$, and into a complete ultrametric

space. Let $K \Subset \Omega$, $m \in \mathbb{N}_0$ and $p \geq 0$. Then $V(K, m, p) \subset \mathcal{G}(\Omega)$ is defined as the collection of elements of $u \in \mathcal{G}(\Omega)$ with a representative u_ε such that

$$\sup_{x \in K} \sup_{|\alpha| \leq m} |\partial^\alpha u_\varepsilon(x)| = \mathcal{O}(\varepsilon^{-p}) \text{ as } \varepsilon \rightarrow 0.$$

The sets $V(K, m, p) \subset \mathcal{G}(\Omega)$ define a base of neighborhoods of zero for a topology with the properties mentioned above. This topology was introduced in [4]; its vast potential was discovered by [39] who coined the term *sharp topology*. Important further developments are due to [2, 10].

We need a couple of further notions from the theory of Colombeau generalized functions. Regularity theory is based on the subalgebra $\mathcal{G}^\infty(\Omega)$ of *regular generalized functions* in $\mathcal{G}(\Omega)$. It is defined by those elements which have a representative satisfying

$$\forall K \Subset \Omega \exists p \geq 0 \forall \alpha \in \mathbb{N}_0^n : \sup_{x \in K} |\partial^\alpha u_\varepsilon(x)| = \mathcal{O}(\varepsilon^{-p}) \text{ as } \varepsilon \rightarrow 0. \quad (7)$$

Observe the change of quantifiers with respect to formula (2); locally, all derivatives of a regular generalized function have the same order of growth in $\varepsilon > 0$. One has that (see [27, Thm. 5.2])

$$\mathcal{G}^\infty(\Omega) \cap \mathcal{D}'(\Omega) = \mathcal{C}^\infty(\Omega).$$

For the purpose of describing the regularity of Colombeau generalized functions, $\mathcal{G}^\infty(\Omega)$ plays the same role as $\mathcal{C}^\infty(\Omega)$ does in the setting of distributions. However, $\mathcal{G}^\infty(\Omega)$ is not invariant under nonlinear maps. For this reason, various subspaces of $\mathcal{G}(\Omega)$ have been introduced to measuring regularity in the nonlinear case (see e.g. [29]). We shall make use of just one of them, based on the notion of subsheaf regularity introduced in [19].

Definition 1. $\mathcal{L}_\mathcal{G}^1(\Omega)$ is the space of the elements of $\mathcal{G}(\Omega)$ with a representative $(u_\varepsilon)_{\varepsilon \in (0,1]}$ such that $\lim_{\varepsilon \rightarrow 0} u_\varepsilon$ exists in $L_{\text{loc}}^1(\Omega)$.

Proposition 2. $f(\mathcal{L}_\mathcal{G}^1(\Omega)) \subset \mathcal{L}_\mathcal{G}^1(\Omega)$ for every smooth function f all whose derivatives grow at most polynomially at infinity and which is Lipschitz continuous with a global Lipschitz constant.

Proof: The polynomial bounds guarantee that $f(u)$ is a well-defined element of $\mathcal{G}(\Omega)$. Assume that u_ε converges to an element $w \in L^1(K)$ on some compact set K . The estimate $|f(u_\varepsilon) - f(w)| \leq \text{Lip}_f |u_\varepsilon - w|$ shows that $f(u_\varepsilon)$ converges to $f(w)$, as desired. $\square$

The kind of regularity that is encapsulated in the subspace above is described by the following rather obvious assertion:

$$\mathcal{L}_\mathcal{G}^1(\Omega) \cap \mathcal{D}'(\Omega) = L_{\text{loc}}^1(\Omega).$$

We end this section by recalling the *association relation* on the Colombeau algebra $\mathcal{G}(\Omega)$. It identifies elements of $\mathcal{G}(\Omega)$ if they coincide in the weak limit. That is, $u, v \in \mathcal{G}(\Omega)$ are called associated, $u \approx v$, if $\lim_{\varepsilon \rightarrow 0} \int (u_\varepsilon(x) - v_\varepsilon(x)) \psi(x) dx = 0$ for all test functions $\psi \in \mathcal{D}(\Omega)$. We shall also say that u is associated with a distribution w if $u_\varepsilon \rightarrow w$ in the sense of distributions as $\varepsilon \rightarrow 0$.

3 Existence/uniqueness of generalized solutions

This section is devoted to solving the semilinear wave equation (1) in the Colombeau algebra $\mathcal{G}(\mathbb{R}^2)$. Recall first that if w is a classical solution of the linear wave equation

$$\begin{aligned} \partial_t^2 w(x, t) - \partial_x^2 w(x, t) &= h(x, t), & (x, t) \in \mathbb{R}^2, \\ w(x, 0) &= a(x), \quad \partial_t w(x, 0) = b(x), & x \in \mathbb{R}, \end{aligned} \quad (8)$$

then it solves the integral equation

$$w(x, t) = \frac{1}{2}(a(x-t) + a(x+t)) + \frac{1}{2} \int_{x-t}^{x+t} b(y) dy + \frac{1}{2} \int_0^t \int_{x-t+s}^{x+t-s} h(y, s) dy ds. \quad (9)$$

Let $K_0 = [-\kappa, \kappa]$ be a compact interval. For $0 \leq T \leq \kappa$, the trapezoidal region K_T is defined by

$$K_T = \{(x, t) \in \mathbb{R}^2 : 0 \leq t \leq T, |x| \leq \kappa - t\}. \quad (10)$$

Using (9), the following estimate is easily deduced ($0 \leq t \leq T \leq \kappa$):

$$\|w\|_{L^\infty(K_T)} \leq \|a\|_{L^\infty(K_0)} + T\|b\|_{L^\infty(K_0)} + T \int_0^T \|h\|_{L^\infty(K_s)} ds. \quad (11)$$

For later reference, we note that estimate (11) holds with the L^1 -norm in place of the L^∞ -norm as well. We now turn to the semilinear wave equation (1). We assume throughout that $u \rightarrow f(u)$ is a smooth function all whose derivatives are of at most polynomial growth as $|u| \rightarrow \infty$, that f satisfies a global Lipschitz estimate (i.e., has a bounded first derivative) and that $f(0) = 0$.

Proposition 3. *Assume that the function f is as described above. Let $a, b \in \mathcal{G}(\mathbb{R})$ and $h \in \mathcal{G}(\mathbb{R}^2)$. Then problem (1) has a unique solution $u \in \mathcal{G}(\mathbb{R}^2)$. The solution depends continuously on the data with respect to the sharp topology.*

Proof: To prove the existence of a solution, take representatives $a_\varepsilon, b_\varepsilon, h_\varepsilon$ of a, b, h , respectively, and let $u_\varepsilon \in \mathcal{C}^\infty(\mathbb{R}^2)$ be the unique classical solution to the semilinear wave equation with regularized data:

$$\begin{aligned} \partial_t^2 u_\varepsilon - \partial_x^2 u_\varepsilon &= f(u_\varepsilon) + h_\varepsilon & \text{on } \mathbb{R}^2, \\ u_\varepsilon(\cdot, 0) &= a_\varepsilon, \quad \partial_t u_\varepsilon(\cdot, 0) = b_\varepsilon & \text{on } \mathbb{R}. \end{aligned} \quad (12)$$

The classical solution u_ε to (12) is constructed by rewriting (12) as an integral equation and invoking a fixed point argument (this involves applying estimate (11) successively to all derivatives). If we show that the net $(u_\varepsilon)_{\varepsilon \in (0, 1]}$ belongs to $\mathcal{E}_M(\mathbb{R}^2)$, its equivalence class in $\mathcal{G}(\mathbb{R}^2)$ will be a solution. We shall do the proof only for the upper half-plane $(x, t) \in \mathbb{R} \times [0, \infty)$; the arguments for $t \leq 0$ are similar. To show that the zero-th derivative of u_ε satisfies the estimate (2), we take a region K_T and invoke inequality (11) to see that

$$\|u_\varepsilon\|_{L^\infty(K_t)} \leq \|a_\varepsilon\|_{L^\infty(K_0)} + T\|b_\varepsilon\|_{L^\infty(K_0)} + T \int_0^t \|f(u_\varepsilon) + h_\varepsilon\|_{L^\infty(K_s)} ds. \quad (13)$$

The last term on the right hand side of (13) is estimated by

$$T \int_0^t \|f'\|_{L^\infty(\mathbb{R})} \|u_\varepsilon\|_{L^\infty(K_s)} ds + T^2 \|h_\varepsilon\|_{L^\infty(K_T)} \quad (14)$$

Using that each of the terms involving $a_\varepsilon, b_\varepsilon, h_\varepsilon$ is of order $\mathcal{O}(\varepsilon^{-p})$ for some p , we infer from Gronwall's inequality that the same is true of $\|u_\varepsilon\|_{L^\infty(K_t)}$ for $0 \leq t \leq T$. Thus u_ε is moderate on the region K_T , that is, it satisfies the estimate (2) there. To get the estimates for the higher order derivatives, one just differentiates the equation and employs the same arguments inductively, using that the lower order terms are already known to be moderate from the previous steps.

To prove uniqueness, we consider representatives $u_\varepsilon, v_\varepsilon \in \mathcal{E}_M[\mathbb{R}^2]$ of two solutions u and v . Their difference satisfies

$$\begin{aligned} \partial_t^2(u_\varepsilon - v_\varepsilon) - \partial_x^2(u_\varepsilon - v_\varepsilon) &= (f(u_\varepsilon) - f(v_\varepsilon)) + n_\varepsilon, \\ (u_\varepsilon(\cdot, 0) - v_\varepsilon(\cdot, 0)) &= n_{0\varepsilon}, \quad \partial_t(u_\varepsilon(\cdot, 0) - v_\varepsilon(\cdot, 0)) = n_{1\varepsilon} \end{aligned}$$

for certain null elements $n_\varepsilon, n_{0\varepsilon}, n_{1\varepsilon}$. Thus $u_\varepsilon - v_\varepsilon$ satisfies an estimate of the form (13), but with the null elements $n_\varepsilon, n_{0\varepsilon}, n_{1\varepsilon}$ replacing $a_\varepsilon, b_\varepsilon, h_\varepsilon$ there. This implies as above that the L^∞ -norm of $u_\varepsilon - v_\varepsilon$ on K_T is of order $\mathcal{O}(\varepsilon^q)$ for every $q \geq 0$. By [14, Thm. 1.2.3], the null estimate (3) on $u_\varepsilon - v_\varepsilon$ suffices to have null estimates on all derivatives. Thus $u = v$ in $\mathcal{G}(\mathbb{R}^2)$.

The proof of continuous dependence on the data a, b, h follows exactly the same lines. $\square$

Having established a general existence- and uniqueness result, the question arises what else can be said about the solution, when more is known about the data. A number of results on the qualitative properties are indeed available. We begin with the most basic result on $\mathcal{C}^\infty$ -smoothness of the solution when the data are smooth.

Proposition 4. *Assume that a, b belong to $\mathcal{C}^\infty(\mathbb{R})$ and h is in $\mathcal{C}^\infty(\mathbb{R}^2)$. Then the generalized solution $u \in \mathcal{G}(\mathbb{R}^2)$ to problem (1) coincides with the classical solution $w \in \mathcal{C}^\infty(\mathbb{R}^2)$, that is, $u = \iota(w)$ in $\mathcal{G}(\mathbb{R}^2)$.*

Proof: The imbeddings ι and σ coincide on $\mathcal{C}^\infty(\mathbb{R})$ and on $\mathcal{C}^\infty(\mathbb{R}^2)$. Thus we may represent the data by $a_\varepsilon \equiv a$, $b_\varepsilon \equiv b$ and $h_\varepsilon \equiv h$. The proof of Prop. 3 shows that $u_\varepsilon \equiv w$ is a representative of the generalized solution, and this means that $u = \iota(w)$ in $\mathcal{G}(\mathbb{R}^2)$. $\square$

Proposition 5. *Assume that the data a, b and h are continuous functions and let $w \in \mathcal{C}(\mathbb{R}^2)$ be the corresponding continuous (weak) solution. Then the generalized solution $u \in \mathcal{G}(\mathbb{R}^2)$ is associated with w , that is, $u \approx \iota(w)$.*

Proof: Let $a_\varepsilon = a * \varphi_\varepsilon$ and similarly for b_ε and h_ε with appropriate one- or two-dimensional mollifiers φ_ε . On the one hand, the families $a_\varepsilon, b_\varepsilon, h_\varepsilon$ define representatives of the generalized functions $\iota(a), \iota(b), \iota(h)$. On the other hand, they converge locally uniformly to the original continuous functions a, b, h . The estimates (13) and (14), applied to $u_\varepsilon - w$, $a_\varepsilon - a$ etc. together with Gronwall's inequality, show that the solution u_ε converge to w with respect the sup-norm on compact sets. In particular, u is associated with w . $\square$

Note that in general $u \neq \iota(w)$ as elements of $\mathcal{G}(\mathbb{R}^{n+1})$. As a rule, equality of continuous and generalized solutions does not hold in any algebra of generalized functions. This is similar to Schwartz' impossibility result and has been worked out in [25]. When the data are distributions, there may be no meaning for a distributional solution, in general. Yet the solution in $\mathcal{G}(\mathbb{R}^2)$ may still be associated with a distribution. We will turn to an incident of such a situation in the next section.

Proposition 3 is a model result. In fact, for vanishing driving term $h \equiv 0$, it is a special case of [3] and of [23]; for Lipschitz continuous, smooth f , existence and uniqueness of a solution in $\mathcal{G}(\mathbb{R}^2)$ can be proven in space dimensions $n = 1, 2, 3$, see e.g. [28]. If f is not Lipschitz, but of polynomial growth, energy estimates can be used to construct solutions in the Colombeau algebra $\mathcal{G}_{2,2}(\mathbb{R}^n \times [0, \infty))$ introduced in [5]. As in the classical case, the growth type of f is connected with the space dimension; the cases $1 \leq n \leq 6$ have been treated in [21]; for $n = 3$ see also [7].

4 Delta waves

In this section we investigate the behavior of the solution $u \in \mathcal{G}(\mathbb{R}^2)$ to the semilinear wave equation (1) when the initial data a, b are given by distributions. In certain cases, the smooth solutions u_ε with regularized initial data $a_\varepsilon, b_\varepsilon$ converge to a distributional limit. When it exists, this limit is called a *delta wave*. Stated equivalently, the generalized solution $u \in \mathcal{G}(\mathbb{R}^2)$ with data $\iota(a), \iota(b)$ admits an associated distribution. To be sure, this distribution is not a solution of the original nonlinear wave equation in any sense. It rather has to be viewed as describing the qualitative properties of the Colombeau solution.

For simplicity of exposition, we will assume here that the right hand side h is zero (see e.g. [28] for a result with nonzero h) and will exhibit the model case of bounded nonlinearity f and initial data with discrete support. Thus $a, b \in \mathcal{D}'(\mathbb{R})$ will be assumed to be Dirac measures or their derivatives situated at a finite set of points. Consider the weak solution $v \in \mathcal{D}'(\mathbb{R}^2)$ to the linear problem

$$\begin{aligned} \partial_t^2 v - \partial_x^2 v &= 0 & \text{on } \mathbb{R}^2, \\ v(\cdot, 0) = a, \partial_t v(\cdot, 0) &= b & \text{on } \mathbb{R}. \end{aligned} \quad (15)$$

By d'Alembert's formula, it is given by

$$v(x, t) = \frac{1}{2}(a(x-t) + a(x+t)) + \frac{1}{2} \int_{x-t}^{x+t} b(y) dy. \quad (16)$$

Its singular support S consists of the characteristic lines with slope ± 1 emanating from the points of support of a, b , hence has Lebesgue measure zero. Denote by $\mathbf{1}_{Sc} v$ the almost everywhere defined, measurable function equal to $v(x, t)$ for $(x, t) \notin S$. In fact, the function $\mathbf{1}_{Sc} v$ is piecewise constant and arises from the measure part in b solely. For example, when $b(x) = \delta(x)$, we have that $\mathbf{1}_{Sc} v(x, t) = 1/2$ for $|x| < t$ and $\mathbf{1}_{Sc} v(x, t) = 0$ otherwise. If the function f is continuous, then the nonlinear equation

$$\begin{aligned} \partial_t^2 w - \partial_x^2 w &= f(\mathbf{1}_{Sc} v + w) & \text{on } \mathbb{R}^2, \\ w(\cdot, 0) = 0, \partial_t w(\cdot, 0) &= 0 & \text{on } \mathbb{R} \end{aligned} \quad (17)$$

has a unique weak solution $w \in \mathcal{C}(\mathbb{R}^2)$. Finally, let $u \in \mathcal{G}(\mathbb{R}^2)$ be the Colombeau solution to the nonlinear problem

$$\begin{aligned} \partial_t^2 u - \partial_x^2 u &= f(u) && \text{on } \mathbb{R}^2, \\ u(\cdot, 0) &= \iota(a), \quad \partial_t u(\cdot, 0) = \iota(b) && \text{on } \mathbb{R}. \end{aligned} \quad (18)$$

As we shall see shortly, $v + w$ is a delta wave, i.e. the associated distribution corresponding to the Colombeau solution to (18).

Proposition 6. *Assume that the data $a, b \in \mathcal{D}'(\mathbb{R})$ have discrete support, f is smooth, bounded and globally Lipschitz. Then the generalized solution $u \in \mathcal{G}(\mathbb{R}^2)$ is associated with the sum of the distributions $v \in \mathcal{D}'(\mathbb{R}^2)$ and the continuous function $w \in \mathcal{C}(\mathbb{R}^2)$, that is, $u \approx \iota(v + w)$.*

Proof: The solution $u \in \mathcal{G}(\mathbb{R}^2)$ has a representative which satisfies

$$\begin{aligned} \partial_t^2 u_\varepsilon - \partial_x^2 u_\varepsilon &= f(u_\varepsilon) && \text{on } \mathbb{R}^2, \\ u_\varepsilon(\cdot, 0) &= a_\varepsilon, \quad \partial_t u_\varepsilon(\cdot, 0) = b_\varepsilon && \text{on } \mathbb{R} \end{aligned}$$

where $a_\varepsilon = a * \varphi_\varepsilon, b_\varepsilon = b * \varphi_\varepsilon$. Further, let v_ε be the classical smooth solution to the linear wave equation (15) with data $a_\varepsilon, b_\varepsilon$. We have

$$\begin{aligned} (\partial_t^2 u - \partial_x^2)(u_\varepsilon - v_\varepsilon - w) &= f(u_\varepsilon) - f(\mathbf{1}_{S^c} v + w) && \text{on } \mathbb{R}^2, \\ (u_\varepsilon - v_\varepsilon - w)(\cdot, 0) &= 0, \quad \partial_t(u_\varepsilon - v_\varepsilon - w)(\cdot, 0) = 0 && \text{on } \mathbb{R}. \end{aligned} \quad (19)$$

We rewrite the right hand side of the first line in (19) as

$$f(u_\varepsilon) - f(v_\varepsilon + w) + f(v_\varepsilon + w) - f(\mathbf{1}_{S^c} v + w).$$

Using (11) with the L^1 -norms, we arrive at

$$\begin{aligned} \|u_\varepsilon - v_\varepsilon - w\|_{L^1(K_t)} &\leq T \int_0^t \|f'\|_{L^\infty(\mathbb{R})} \|u_\varepsilon - v_\varepsilon - w\|_{L^1(K_s)} ds \\ &+ T^2 \|f(v_\varepsilon + w) - f(\mathbf{1}_{S^c} v + w)\|_{L^1(K_T)} \end{aligned}$$

valid for $0 \leq t \leq T$. Since $f(v_\varepsilon + w) - f(\mathbf{1}_{S^c} v + w)$ converges to zero almost everywhere and remains bounded, Lebesgue's theorem shows that its L^1 -norm on K_T converges to zero. By Gronwall's lemma, it follows that the L^1 -norm of $u_\varepsilon - v_\varepsilon - w$ converges to zero on any K_T as well. Hence u_ε converges to $v + w$ weakly, which translates into the claimed association result. $\square$

This result is an example of a wide range of much more general results on the existence of delta waves. We refer to the papers [22, 26, 33, 37]. In case the driving term h is white noise, the generalized solution to the semilinear (stochastic) wave equation is associated with the solution of a linear stochastic wave equation in many cases. This has been shown e.g. in [1, 30, 32]. White noise or positive noise in the initial data has been considered in [31, 35].

5 Linear regularity theory

Consider the linear wave equation (15) with initial data distributions $a, b \in \mathcal{D}'(\mathbb{R})$. The classical propagation of singularities results says that the singular

support of the solution $v \in \mathcal{D}'(\mathbb{R}^2)$ is contained in the union of characteristic lines emanating from the singular support of the initial data. For example, when the singular support of the initial data is situated at the origin, the solution is a smooth function outside the forward and backward light-cone $S = \{(x, t) \in \mathbb{R}^2 : |x| = |t|\}$. In the Colombeau setting, this is not so, as shown by the following example.

Example 7. Let $u \in \mathcal{G}(\mathbb{R}^2)$ be the Colombeau solution to the linear wave equation

$$\begin{aligned} \partial_t^2 u - \partial_x^2 u &= 0 & \text{on } \mathbb{R}^2, \\ u(\cdot, 0) = 0, \partial_t u(\cdot, 0) &= \iota(\delta)^2 & \text{on } \mathbb{R}. \end{aligned}$$

with initial data given by the square of the Dirac measure in $\mathcal{G}(\mathbb{R})$. Then u does not coincide with a $\mathcal{C}^\infty$ -function in the interior of the light-cone $\{(x, t) \in \mathbb{R}^2 : |x| < |t|\}$.

In fact, by d'Alembert's formula, the solution $u \in \mathcal{G}(\mathbb{R}^2)$ has the value

$$\frac{1}{2} \int_{-\infty}^{\infty} \iota(\delta)^2(x) dx = \text{class of } \left(\frac{1}{2} \int_{-\infty}^{\infty} \varphi_\varepsilon^2(x) dx \right)_{\varepsilon \in (0,1]}$$

inside the forward light-cone $\{(x, t) \in \mathbb{R}^2 : |x| < t\}$, $t > 0$, which is a generalized constant not belonging to $\mathbb{C}$.

The example shows that the space $\mathcal{C}^\infty$ is not suitable for measuring regularity in Colombeau algebras. Its role is taken over by the algebra $\mathcal{G}^\infty$ defined in the Introduction. We have the following result.

Proposition 8. Let $v \in \mathcal{G}(\mathbb{R}^2)$ be the Colombeau solution to the linear wave equation (15) with initial data $a, b \in \mathcal{G}(\mathbb{R})$. If $a, b \in \mathcal{G}^\infty(\mathbb{R} \setminus \{0\})$, then $u \in \mathcal{G}^\infty(\mathbb{R} \setminus S)$.

Proof: We take a cut-off function $\chi \in \mathcal{C}^\infty(\mathbb{R})$ which is equal to one in a small neighborhood of the origin and zero otherwise. Then $v = v_1 + v_2$ where $v_1 \in \mathcal{G}(\mathbb{R}^2)$ is the solution with initial data $a_1 = (1-\chi)a$, $b_1 = (1-\chi)b$ and $v_2 \in \mathcal{G}(\mathbb{R}^2)$ is the solution with initial data $a_2 = \chi a$, $b_2 = \chi b$. Using d'Alembert's formula, we have that

$$\partial_x v_{1\varepsilon} = \frac{1}{2} (a'_{1\varepsilon}(x+t) + a'_{1\varepsilon}(x-t) + b_{1\varepsilon}(x+t) - b_{1\varepsilon}(x-t))$$

and similarly for $\partial_t v_{1\varepsilon}$. Thus if all the derivatives of a_ε and b_ε are of the same local order, so are the derivatives of $v_{1\varepsilon}$. This shows that v_1 belongs to $\mathcal{G}^\infty(\mathbb{R}^2)$. Similarly, differentiating v_2 once shows that the first and higher derivatives of v_2 vanish outside a small neighborhood of S , determined by the diameter of the support of χ . Since χ was arbitrary, this shows that v_2 is a generalized constant on $\mathbb{R} \setminus S$, thus belongs to $\mathcal{G}^\infty(\mathbb{R} \setminus S)$ as well. $\square$

It is clear that Prop. 8 can be generalized to initial data with discrete $\mathcal{G}^\infty$ -singular support. It also holds in any space dimension [27].

6 Nonlinear regularity theory

This section addresses propagation of singularities for the semilinear wave equation. The initial data that are assumed to be regular outside the origin, but

may have a singularity there. Following the classical theory of propagation of jump discontinuities one may hope to prove that, in the one-dimensional case, the solution will be regular (in the appropriate sense) in the interior of the light-cone S . That this behavior occurs in the $\mathcal{C}^\infty$ -category for the semilinear wave equation in one space dimension was shown by [38]; it does not hold in higher space dimensions or for higher order operators [36], for which *anomalous* singularities may occur. The question we ask is whether this transport of regularity into the interior of the light-cone happens in the Colombeau setting as well. We do not have a general result yet, but we will be able to prove this transport of regularity under conditions which resemble the situation leading to *delta waves*. We shall employ the subalgebra $\mathcal{G}^\infty$ and the space $\mathcal{L}_{\mathcal{G}}^1$ defined in the Introduction for measuring the regularity.

We consider the semilinear wave equation (1). As before, we assume that the function $u \rightarrow f(u)$ is smooth with all derivatives of at most polynomial growth as $|u| \rightarrow \infty$, and that it satisfies a global Lipschitz estimate, i.e., has a globally bounded derivative. We take initial data of the form

$$u_i = r_i + s_i \in \mathcal{G}(\mathbb{R}), \quad i = 0, 1$$

where

$$r_i \in \mathcal{L}_{\mathcal{G}}^1(\mathbb{R}) \text{ and } \text{supp}(s_i) = \{0\}, \quad i = 0, 1.$$

Define the generalized complex number $M \in \tilde{\mathbb{C}}$ as the class of

$$M_\varepsilon = \frac{1}{2} \int_{-\infty}^{\infty} s_{1\varepsilon}(x) dx$$

where $(s_{1\varepsilon})_{\varepsilon \in (0,1]}$ is a representative of s_1 . We shall write $M \approx m$ if M_ε converges to a complex number $m \in \mathbb{C}$ and $|M| \approx \infty$ if $|M_\varepsilon| \rightarrow \infty$ as $\varepsilon \rightarrow 0$. Again $S = \{(x, t) \in \mathbb{R}^2 : |x| = |t|\}$ denotes the forward and backward light-cone.

Proposition 9. *Assume that the function f is smooth, globally Lipschitz, and all its derivatives are polynomially bounded. Let $u \in \mathcal{G}(\mathbb{R}^2)$ be the solution to the semilinear wave equation (1) with initial data u_0, u_1 as described above. If either*

- (a) *f is globally bounded and $M \approx m$ for some $m \in \mathbb{C}$, or*
- (b) *f is globally bounded, $\lim_{|y| \rightarrow \infty} f(y)$ exists and $|M| \approx \infty$*

then $u \in \mathcal{G}^\infty(\mathbb{R}^2 \setminus S) + \mathcal{L}_{\mathcal{G}}^1(\mathbb{R}^2)$.

Proof: We first derive the estimates for (x, t) in the upper half-plane $\mathbb{R} \times [0, \infty)$. Denote by $\mathbf{1}_\Sigma$ the characteristic function of the solid light-cone $\Sigma = \{(x, t) : t \geq 0, |x| \leq t\}$. Let $(r_{i\varepsilon})_{\varepsilon \in (0,1]}$, $(s_{i\varepsilon})_{\varepsilon \in (0,1]}$, $i = 0, 1$, be representatives of the initial data, where we may assume that $\text{supp}(s_{i\varepsilon}) \subset [-\eta, \eta]$ with η as small as we wish. Let $u_\varepsilon, v_\varepsilon \in \mathcal{C}^\infty(\mathbb{R} \times [0, \infty))$, $w_\varepsilon \in L_{\text{loc}}^1(\mathbb{R} \times [0, \infty))$ be the solutions to

$$\begin{aligned} (\partial_t^2 - \partial_x^2) u_\varepsilon &= f(u_\varepsilon), & u_\varepsilon(\cdot, 0) &= u_{0\varepsilon}, & \partial_t u_\varepsilon(\cdot, 0) &= u_{1\varepsilon}, \\ (\partial_t^2 - \partial_x^2) v_\varepsilon &= 0, & v_\varepsilon(\cdot, 0) &= s_{0\varepsilon}, & \partial_t v_\varepsilon(\cdot, 0) &= s_{1\varepsilon}, \\ (\partial_t^2 - \partial_x^2) w_\varepsilon &= f(M_\varepsilon \mathbf{1}_\Sigma + w_\varepsilon), & w_\varepsilon(\cdot, 0) &= r_{0\varepsilon}, & \partial_t w_\varepsilon(\cdot, 0) &= r_{1\varepsilon}. \end{aligned}$$

Then

$$(\partial_t^2 - \partial_x^2) (u_\varepsilon - v_\varepsilon - w_\varepsilon) = f(u_\varepsilon) - f(v_\varepsilon + w_\varepsilon) + f(v_\varepsilon + w_\varepsilon) - f(M_\varepsilon \mathbf{1}_\Sigma + w_\varepsilon)$$

with zero initial data. The difference of the first two terms on the right hand side can be estimated by the L^∞ -norm of f' times $|u_\varepsilon - v_\varepsilon - w_\varepsilon|$, while the difference of the last two terms vanishes off an η -neighborhood of the light-cone S , because $v_\varepsilon(x, t) = M_\varepsilon$ for $|x| < t - \eta$. Taking a trapezoidal region K_T as in (10), the boundedness of f , inequality (11) and Gronwall's lemma give an estimate of the form

$$\|u_\varepsilon - v_\varepsilon - w_\varepsilon\|_{L^1(K_T)} \leq C\eta \quad (20)$$

for some constant $C > 0$ and all $\varepsilon \in (0, 1]$. In the case (a), let $w \in L^1_{\text{loc}}(\mathbb{R} \times [0, \infty))$ be the solution to

$$(\partial_t^2 - \partial_x^2) w = f(m\mathbf{1}_\Sigma + w), \quad w(\cdot, 0) = r_0, \quad \partial_t w(\cdot, 0) = r_1$$

where r_i is the limit in $L^1_{\text{loc}}(\mathbb{R})$ of $r_{i\varepsilon}$ as $\varepsilon \rightarrow 0$, $i = 0, 1$. Now

$$(\partial_t^2 - \partial_x^2)(w_\varepsilon - w) = f(M_\varepsilon\mathbf{1}_\Sigma + w_\varepsilon) - f(M_\varepsilon\mathbf{1}_\Sigma + w) + f(M_\varepsilon\mathbf{1}_\Sigma + w) - f(m\mathbf{1}_\Sigma + w)$$

with initial data given by $r_{i\varepsilon} - r_i$, $i = 0, 1$. The difference of the first two terms on the right hand side is bounded by the L^∞ -norm of f' times $|w_\varepsilon - w|$. By assumption, the difference of the last two terms converges to zero almost everywhere. Using inequality (11) with L^1 -norms in place of the L^∞ -norms and Gronwall's lemma as above shows that

$$\|w_\varepsilon - w\|_{L^1(K_T)} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0. \quad (21)$$

In case (b), defining $w \in L^1_{\text{loc}}(\mathbb{R} \times [0, \infty))$ as the solution to

$$(\partial_t^2 - \partial_x^2) w = L\mathbf{1}_\Sigma, \quad w(\cdot, 0) = r_0, \quad \partial_t w(\cdot, 0) = r_1,$$

the same argument as above leads to the convergence result (21) in this case as well. Combining (20) which holds for arbitrarily chosen $\eta > 0$ with (21) shows that $u_\varepsilon - v_\varepsilon$ converges to w in $L^1(K_T)$ as $\varepsilon \rightarrow 0$. By Prop. 8, v_ε enjoys the $\mathcal{G}^\infty$ -estimate (7) off the light-cone S . Thus $u_\varepsilon = v_\varepsilon + (u_\varepsilon - v_\varepsilon)$ defines an element of $\mathcal{G}^\infty((\mathbb{R} \times [0, \infty)) \setminus S) + \mathcal{L}^1_{\mathcal{G}}(\mathbb{R} \times [0, \infty))$.

The arguments for (x, t) in the lower half-plane $\mathbb{R} \times (-\infty, 0]$ are the same with $-M$ in place of M . $\square$

Remark 10. *The hypotheses on the generalized constant M in Prop. 9 are satisfied when the term s_1 in the initial data is a polynomial in the Dirac measure and its derivatives. In fact, when $s_1 = \partial^\alpha \iota(\delta)$, we have $M \approx 1$ for $\alpha = 0$ and $M \approx 0$ for $\alpha > 0$. When $s_1 = \pi(\iota(\delta))$ for some polynomial function π , only the cases $M \approx m$ for some $m \in \mathbb{C}$ or $|M| \approx \infty$ can occur.*

Prop. 9 shows that regularity of the type $\mathcal{G}^\infty + \mathcal{L}^1_{\mathcal{G}}$ is propagated into the region inside the light-cone. It is clear that it can be generalized in various ways: for example, the support of the singular part s_0, s_1 of the data could consist of a discrete set rather than just a point. However, it should be noted that continuous dependence of the regularized solutions on the data in terms the L^1 -norm enters into the argument, and such a property depends decisively on the particular equation, the space dimension and the nonlinearity f . Prop. 9 exploits such special properties and thus falls short of providing a prototypical description of nonlinear propagation of regularity for Colombeau solutions, which remains a challenging open question.

Another promising approach to measuring regularity in the nonlinear case is the Colombeau-Hölder-Zygmund-scale which has been introduced and applied to nonlinear scalar first order equations in [16].

References

- [1] S. Albeverio, Z. Haba, F. Russo, Trivial solutions for a non-linear two-space dimensional wave equation perturbed by space-time white noise. *Stochastics and Stochastic Reports* **56**(1996), 127 - 160.
- [2] J. Aragona, S. Juriaans, Some structural properties of the topological ring of Colombeau's generalized numbers. *Communications in Algebra* **29**(2001), 2201 - 2230.
- [3] H. A. Biagioni, The Cauchy problem for semilinear hyperbolic systems with generalized functions as initial conditions. *Result. Math.* **14**(1988), 232 - 241.
- [4] H. A. Biagioni. *A Nonlinear Theory of Generalized Functions*. Lect. Notes Math. 1421. Springer-Verlag, Berlin, 1990.
- [5] H. A. Biagioni. M. Oberguggenberger, Generalized solutions to the Korteweg - de Vries and the regularized long-wave equations. *SIAM J. Math. Anal.* **23**(1992), 923 - 940.
- [6] J. F. Colombeau. *New Generalized Functions and Multiplication of Distributions*. North-Holland Math. Studies 84. North-Holland, Amsterdam, 1984.
- [7] J. F. Colombeau. *Elementary Introduction to New Generalized Functions*. North-Holland Math. Studies 113. North-Holland, Amsterdam, 1985.
- [8] J. F. Colombeau, M. Oberguggenberger, On a hyperbolic system with a compatible quadratic term: Generalized solutions, delta waves, and multiplication of distributions. *Comm. Part. Diff. Eqs.* **15**(1990), 905 - 938.
- [9] C. Garetto, Pseudo-differential operators in algebras of generalized functions and global hypoellipticity. *Acta Appl. Math.* **80**(2004), 123 - 174.
- [10] C. Garetto. Topological structures in Colombeau algebras: topological $\tilde{\mathbb{C}}$ -modules and duality theory. *Acta Appl. Math.* **88**(2005), 81 - 123.
- [11] C. Garetto, T. Gramchev, M. Oberguggenberger, Pseudo-differential operators with generalized symbols and regularity theory. *Electron. J. Diff. Eqns.* **2005**(116)(2005), 1 - 43.
- [12] C. Garetto, G. Hörmann, Microlocal analysis of generalized functions: pseudodifferential techniques and propagation of singularities. *Proc. Edinburgh. Math. Soc.* **48**(2005), 603 - 629.
- [13] T. Gramchev. Semilinear hyperbolic systems with singular initial data. *Monatsh. Math.* **112** (1991), 99 - 113.

- [14] M. Grosser, M. Kunzinger, M. Oberguggenberger, R. Steinbauer, *Geometric Theory of Generalized Functions with Applications to General Relativity*. Mathematics and its Applications 537. Kluwer Acad. Publ., Dordrecht, 2001.
- [15] G. Hörmann, First-order hyperbolic pseudodifferential equations with generalized symbols. *J. Math. Anal. Appl.* **293**(2004), 40 - 56.
- [16] G. Hörmann, Hölder-Zygmund regularity in algebras of generalized functions. *Zeitschr. Anal. Anw.* **23**(2004), 139 - 165.
- [17] G. Hörmann, M. V. de Hoop, Microlocal analysis and global solutions of some hyperbolic equations with discontinuous coefficients. *Acta Appl. Math.* **67**(2001), 173 - 224.
- [18] F. Lafon, M. Oberguggenberger, Generalized solutions to symmetric hyperbolic systems with discontinuous coefficients: the multidimensional case. *J. Math. Anal. Appl.* **160**(1991), 93 - 106.
- [19] J.-A. Marti. $(\mathcal{C}, \mathcal{E}, \mathcal{P})$ -Sheaf Structures and Applications. In: M. Grosser, G. Hörmann, M. Kunzinger, and M. Oberguggenberger, editors. *Nonlinear Theories of Generalized Functions, Proceedings of the workshop at the Erwin Schrödinger Institute for Mathematical Physics, Vienna 1997*, Boca Raton, 1999. Chapman & Hall/CRC, 175–186.
- [20] M. Nedeljkov, S. Pilipović, D. Scarpalézos. *The Linear Theory of Colombeau Generalized Functions*. Pitman Research Notes in Math. 385. Longman Scientific & Technical, Harlow, 1998.
- [21] M. Nedeljkov, M. Oberguggenberger, S. Pilipović, Generalized solutions to a semilinear wave equation. *Nonlinear Analysis* **61**(2005), 461 - 475.
- [22] M. Oberguggenberger, Weak limits of solutions to semilinear hyperbolic systems. *Math. Ann.* **274**(1986), 599 - 607.
- [23] M. Oberguggenberger, Generalized solutions to semilinear hyperbolic systems. *Monatsh. Math.* **103**(1987), 133 - 144.
- [24] M. Oberguggenberger, Hyperbolic systems with discontinuous coefficients: Generalized solutions and a transmission problem in acoustics. *J. Math. Anal. Appl.* **142**(1989), 452 - 467.
- [25] M. Oberguggenberger, Semilinear wave equations with rough initial data: generalized solutions. In: P. Antosik, A. Kaminski (Eds.), *Generalized Functions and Convergence*. World Sci. Publ., London 1990, 181 - 203.
- [26] M. Oberguggenberger. The Carleman system with positive measures as initial data. *Transport Theory Statist. Phys.* **20** (1991), 177 - 197.
- [27] M. Oberguggenberger, *Multiplication of Distributions and Applications to Partial Differential Equations*. Pitman Research Notes Math. 259, Longman Scientific & Technical, Harlow 1992.
- [28] M. Oberguggenberger, Generalized functions in nonlinear models – a survey. *Nonlinear Analysis* **47/8**(2001), 5029 - 5040.

- [29] M. Oberguggenberger, Regularity theory in Colombeau algebras. Bull. T. CXXXIII Acad. Serbe Sci. Arts, Cl. Sci. Math. Nat., Sci. Math. **31**(2006), 147 - 162.
- [30] M. Oberguggenberger, F. Russo, Nonlinear SPDEs: Colombeau solutions and pathwise limits. In: L. Decreusefond, J. Gjerde, B. Øksendal, A.S. Üstünel (Eds.), *Stochastic Analysis and Related Topics VI*. Birkhäuser, Boston 1998, 319 - 332.
- [31] M. Oberguggenberger, F. Russo, Nonlinear stochastic wave equations. Integral Transf. Special Funct. **6**(1998), 71 - 83.
- [32] M. Oberguggenberger, F. Russo. Singular limiting behavior in nonlinear stochastic wave equations. In: A. B. Cruzeiro, J.-C. Zambrini (Eds.), *Stochastic Analysis and Mathematical Physics*, Birkhäuser, Basel 2001, 87 - 99.
- [33] M. Oberguggenberger, Y.-G. Wang, Delta-waves for semilinear hyperbolic Cauchy problems. Math. Nachr. **166**(1994), 317 - 327.
- [34] M. Oberguggenberger, Y.-G. Wang, Reflection of delta-waves for nonlinear wave equations in one space variable. Nonlinear Analysis **22**(1994), 983 - 992.
- [35] D. Rajter-Ćirić, One-dimensional nonlinear stochastic wave equation and triviality effect. Stoch. Anal. Appl. **23**(2005), 1149 - 1164.
- [36] J. Rauch. Singularities of solutions to semilinear wave equations. J. Math. Pures Appl., IX. Sér. **58**(1979), 299 - 308.
- [37] J. Rauch, M. Reed, Nonlinear superposition and absorption of delta waves in one space dimension. Funct. Anal. **73**(1987), 152 - 178.
- [38] M. Reed. Propagation of singularities for non-linear wave equations in one dimension. Commun. Partial Differ. Equations **3**(1978), 153 - 199.
- [39] D. Scarpalézos. Colombeau's generalized functions: topological structures; microlocal properties. A simplified point of view. Bull. T. CXXI Acad. Serbe Sci. Arts, Cl. Sci. Math. Nat., Sci. Math. **25**(2000), 89 - 114.